arXiv:quant-ph/0112018v1 4 Dec 2001

Continuous variable teleportation of single photon states

Toshiki Ide, Takayoshi Kobayashi

Department of Physics, Faculty of Science, University of Tokyo,

7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan *

Holger F. Hofmann

CREST, Japan Science and Technology Corporation (JST),

Research Institute for Electronic Science,

Hokkaido University,

Sapporo 060-0812, Japan

Akira Furusawa

Department of Applied Physics, Faculty of Engeneering, University of Tokyo,

7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan

(Dated: August 27, 2001)

Abstract

We investigate the changes to a single photon state caused by the non-maximal entanglement in continuous variable quantum teleportation. It is shown that the teleportation measurement introduces field coherence in the output.

PACS numbers: 03.67.-a, 03.67.Hk, 42.50.-p

*Electronic address: ide@femto.phys.s.u-tokyo.ac.jp



FIG. 1: Schematic representation of the quantum teleportation setup.

I. INTRODUCTION

Quantum teleportation is a method for Alice (sender) to transmit an unknown quantum input state to Bob (receiver) at a distant place by sending only classical information using a shared entangled state as a resource [1]. In continuous variable quantum teleportation [2, 3, 4], the available entanglement is non-maximal, limited by the amount of squeezing achieved. Fig.1 shows the setup of a continuous variable quantum teleportation. Alice transmits an unknown quantum state $\mid \psi\rangle_A$ to Bob. Alice and Bob share EPR beams in advance. Alice mixes her input state with the reference EPR beam by a 50% beam-splitter and performs an entanglement measurement of the complex field value β. After Bob gets the information of the field measurement value β from Alice, Bob applies a displacement to the output state by mixing the coherent field of a local oscillator with the output EPR beam B.

As has been shown previously [6], the properties of this transfer process can be summarized by the transfer operator $\hat{T}_q(\beta)$ which describes both the probability distribution $P(\beta)$

of measurement results β and the normalized conditional output state $\mid \psi_{\text{out}}(\beta)\rangle$ for any input state $\mid \psi_{\text{in}}\rangle$, such that

$$\sqrt{P(\beta)} \mid \psi_{\text{out}}(\beta)\rangle = \hat{T}_q(\beta) \mid \psi_{\text{in}}\rangle. \tag{1}$$

In its diagonalized form, this transfer operator reads

$$\hat{T}_q(\beta) = \sqrt{\frac{1-q^2}{\pi}} \sum_{n=0}^{\infty} q^n \hat{D}(\beta) \mid n\rangle_{BA}\langle n \mid \hat{D}(-\beta). \tag{2}$$

The non-maximal entanglement is described by the parameter q, which is 0 for a non-entangled vacuum and 1 for maximal entanglement. In the following, this operator will be applied to characterize the teleportation of a single photon input state, with special consideration of the field coherence created in the output by the teleportation process.

II. TELEPORTATION OF A SINGLE PHOTON STATE

The output of a one photon input state is characterized by

$$\hat{T}_q(\beta) \mid 1\rangle = \sqrt{\frac{1-q^2}{\pi}} e^{-(1-q^2)\frac{|\beta|^2}{2}} \hat{D}((1-q)\beta)((1-q^2)\beta^* \mid 0\rangle + q \mid 1\rangle). \tag{3}$$

The normalized output state is then given by

$$\mid \psi_{\text{out}}(\beta)\rangle \;=\; \frac{1}{\sqrt{q^2 + (1-q^2)^2|\beta|^2}} \hat{D}((1-q)\beta)((1-q^2)\beta^* \mid 0\rangle + q \mid 1\rangle). \tag{4}$$

The output state can be described by a displaced coherent superposition of a zero photon and a one photon component. Both the displacement and the coherence depend on the complex measurement value β. They therefore represent a measurement induced coherence of the output state. We characterize this coherence by the expectation value of the complex field amplitude,

$$\begin{aligned} C_q(\beta) &= \langle \psi_{\text{out}}(\beta) \mid \hat{a} \mid \psi_{\text{out}}(\beta)\rangle \\ &= \frac{q(1-q^2)}{q^2 + (1-q^2)^2|\beta|^2}\beta + (1-q)\beta. \end{aligned} \tag{5}$$

Fig.2 illustrates this dependence of C_q on β for $q = 1/2$. For low values of β, C_q rises sharply, levelling off around $|\beta| \approx 0.5$. At higher values of $|\beta|$, C_q slowly approaches $(1-q)\beta$, as indicated by the dotted line. Since the input photon number state has a field expectation



FIG. 2: The absolute value of β dependence of the field amplitude $C_q^{(1)}(\beta)$ in the case of $q = 0.5$. The curve is approaching $y = 0.5q$.

value of zero, the field coherence in the output is a consequence of the measurement β. As has been argued elsewere [6], β corresponds to the result of a field measurement performed on the input state. This measurement creates coherence by projection onto displaced photon number states as indicated by the diagonalized form of $\hat{T}_q(\beta)$ in equation (2). Since the photon is the quantum mechanical equivalent of field intensity, each photon represents an addition of one quantum unit to the field fluctuations. The measurement then converts the field fluctuation into an actual field. This process is responsible for the rapid increase in coherence at low values of β.

III. CONCLUSIONS

We have investigated the effects of continous variable quantum teleportation on a single photon input state. Because of the non-maximal entanglement used in the teleportation, the measurement of β introduces coherence into the output state. We have quantified this coherence, tracing its origin to the field fluctuations of the single photon input.

---

[1] C.H. Bennet, G. Brassard, C. Crepeau, R.Jozsa, A. Peres, and W.K. Wootters, *Phys. Rev. Lett.* **70**, 1895 (1993).

[2] L. Vaidman, *Phys. Rev. A* **49**, 1473 (1994).

[3] S.L. Braunstein and H.J. Kimble, Phys. Rev. Lett. **80**, 869 (1998).

[4] A. Furusawa, J. L. Sørensen, S. L. Braunstein, C. A. Fuchs, H. J. Kimble, and E. J. Polzik, *Science* **282**, 706 (1998).

[5] S. J. van Enk, *Phys. Rev. A* **60**, 5095 (1999).

[6] H. F. Hofmann, T. Ide, T. Kobayashi, and A. Furusawa, *Phys. Rev. A* **62**, 062304 (2000).